

NEWS RELEASE

Contacts: Geoff Schladow, UC Davis, 530-752-3942

Dennis J. Oliver, Tahoe Regional Planning Agency, 775-450-7311

For Release Immediately

March 19, 2009

LAKE TAHOE CLARITY HOLDS STEADY IN 2008, LATEST ANALYSIS SHOWS

LAKE TAHOE – The waters of Lake Tahoe were clear to an average depth of 69.6 feet in 2008, according to UC Davis scientists who have monitored the lake since 1968. That keeps the clarity measurement in the range where it has been for about the past eight years.

When measurements began in 1968, a white "Secchi disk" lowered into the lake was visible at an average depth of 102.4 feet.

Last year UC Davis reported that data since 2001 suggested water clarity loss at the lake had slowed. That encouraging finding is supported by the 2008 average, which is nearly identical to the 2007 average of 70.1 feet.

Individual clarity measurements in 2008 ranged from a maximum depth of 122.2 feet on April 24 to a minimum of only 36.9 feet on Aug. 5.

In fact, the lake was much less clear than normal during the entire period from mid-July to mid-August, possibly the result of smoke from California wildfires, said John Reuter, associate director of the Tahoe Environmental Research Center. For example, the typical Secchi depth in that summer period from 2003-2007 was 50 feet to 65 feet. But in 2008, the depth was 36.9 feet to 46.8 feet.

The overall summer (June-September) clarity depth was also very shallow at 50.5 feet – the shallowest measured since monitoring began in 1968.

Smoke from wildfires elsewhere in California drifted into the Tahoe Basin and lingered for weeks last summer. Prolonged fallout of ash particles could have both blocked light



penetration into the water and fed the lake's algae, which absorb sunlight and reduce water clarity.

According to Geoff Schladow, director of the UC Davis Tahoe Environmental Research Center, "What 2008 highlighted is the temporary impact that wildfires and other factors outside our direct control can have on Lake Tahoe. While progress is being made in both understanding and addressing the root causes of clarity decline, the path to achieving the desired clarity will not be a straight one."

UC Davis and many other academic institutions and public agencies are working together with the private sector to restore and preserve the Tahoe Basin ecosystem. Led by the Tahoe Regional Planning Agency, the collaborative Environmental Improvement Program is a public-private partnership rivaling the most ambitious of US restoration initiatives in its scope.

"We are encouraged that the long-term trend showing clarity loss slowing has held," said Joanne Marchetta, TRPA Executive Director. "Despite the apparent impact seen from wildfires outside the Basin last year, Lake Tahoe's future appears hopeful."

UC Davis researchers measure the lake's clarity with a Secchi disk every seven to 10 days at two fixed locations. The depth at which the white disk, the size of a dinner plate, disappears from sight is referred to as the Secchi depth.

Their long-term research strongly indicates that Lake Tahoe's dwindling clarity is caused by fine particles and nutrients in the lake. The particles and nutrients enter the lake through erosion, runoff and atmospheric deposition. Once in the lake, they affect clarity by scattering light and by fueling the growth of algae, which absorb light.

"With our Environmental Improvement Program and other initiatives, we feel we may have turned the corner in the effort to restore lake clarity to levels seen a generation ago," said Marchetta. "With a continued commitment to the lake from all sectors, we will accomplish our goal of preserving and protecting Lake Tahoe."

For more information:

-- www.terc.ucdavis.edu or www.trpa.org

Media contacts:

Geoff Schladow, Tahoe Environmental Research Center, (530) 902-2272, gschladow@ucdavis.edu; John Reuter, Tahoe Environmental Research Center, (530) 304-1473 jereuter@ucdavis.edu; Sylvia Wright, UC Davis News Service, (530) 752-7704, swright@ucdavis.edu; Dennis J. Oliver, Tahoe Regional Planning Agency, (775) 589-5235, doliver@trpa.org

Historical Clarity Data -- The annual average Secchi depth measurements for the past several years were as follows:

- * 2008: 69.6 feet (21.2 meters)
- * 2007: 70.2 feet (21.4 meters)
- * 2006: 67.7 feet (20.6 meters)
- * 2005: 72.4 feet (22.1 meters)
- * 2004: 73.6 feet (22.4 meters)
- * 2003: 71 feet (21.6 meters)
- * 2002: 78 feet (23.8 meters)
- * 2001: 73.6 feet (22.4 meters)
- * 2000: 67.3 feet (20.5 meters)
- * 1999: 69 feet (21 meters)
- * 1998: 66 feet (20.1 meters)
- * 1997: 64 feet (19.5 meters)

###